



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Acupuncture for Chronic Pain: Update of an Individual Patient Data Meta-Analysis

Vickers, Andrew J ; Vertosick, Emily A ; Lewith, George ; MacPherson, Hugh ; Foster, Nadine E ;
Sherman, Karen J ; Irnich, Dominik ; Witt, Claudia M ; Linde, Klaus ; Acupuncture Trialists'
Collaboration

DOI: <https://doi.org/10.1016/j.jpain.2017.11.005>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-144283>

Journal Article

Accepted Version

Originally published at:

Vickers, Andrew J; Vertosick, Emily A; Lewith, George; MacPherson, Hugh; Foster, Nadine E; Sherman, Karen J; Irnich, Dominik; Witt, Claudia M; Linde, Klaus; Acupuncture Trialists' Collaboration (2018). Acupuncture for Chronic Pain: Update of an Individual Patient Data Meta-Analysis. *Journal of Pain*, 19(5):455-474.

DOI: <https://doi.org/10.1016/j.jpain.2017.11.005>


Accepted Manuscript

Title: Acupuncture for Chronic Pain: Update of an Individual Patient Data Meta-Analysis

Author: Andrew J. Vickers, Emily A. Vertosick, George Lewith, Hugh MacPherson, Nadine E. Foster, Karen J. Sherman, Dominik Irnich, Claudia M. Witt, Klaus Linde, on behalf of the Acupuncture Trialists' Collaboration

PII: S1526-5900(17)30780-0
DOI: <https://doi.org/10.1016/j.jpain.2017.11.005>
Reference: YJPAI 3487

To appear in: *The Journal of Pain*



Please cite this article as: Andrew J. Vickers, Emily A. Vertosick, George Lewith, Hugh MacPherson, Nadine E. Foster, Karen J. Sherman, Dominik Irnich, Claudia M. Witt, Klaus Linde, on behalf of the Acupuncture Trialists' Collaboration, Acupuncture for Chronic Pain: Update of an Individual Patient Data Meta-Analysis, *The Journal of Pain* (2017), <https://doi.org/10.1016/j.jpain.2017.11.005>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Acupuncture for chronic pain: update of an individual patient data meta-analysis

Andrew J. Vickers PhD, Emily A. Vertosick MPH, George Lewith MD, Hugh MacPherson PhD, Nadine E. Foster DPhil, Karen J. Sherman PhD, Dominik Irnich MD, Claudia M. Witt MD, Klaus Linde MD on behalf of the Acupuncture Trialists' Collaboration

Andrew J. Vickers, DPhil, Memorial Sloan Kettering Cancer Center, New York, NY

Emily A. Vertosick, MPH, Memorial Sloan Kettering Cancer Center, New York, NY

George Lewith, MD, University of Southampton, Southampton, UK

Hugh MacPherson, PhD, University of York, York, UK

Nadine E. Foster, DPhil, Keele University, Newcastle-under-Lyme, UK

Karen J. Sherman, PhD, Group Health Research Institute, Seattle, WA

Dominik Irnich, MD, Ludwig-Maximilians-Universität München (LMU Munich), Munich, Germany

Claudia M. Witt, MD, University of Zurich and University Hospital Zurich, Zurich, Switzerland; Charité-Universitätsmedizin, Berlin, Germany; University of Maryland School of Medicine, Baltimore, Maryland

Klaus Linde, MD, Technical University Munich, Germany

Corresponding author

Andrew J. Vickers
Memorial Sloan Kettering Cancer Center
485 Lexington Avenue
New York, NY 10017 USA
Email: vickersa@mskcc.org
(P): 646-888-8233

Authors' contributions

The study was conceived by AV, GL, CW and KL. AV was responsible for the overall study design; EV for the systematic review; GL and HM with respect to acupuncture analyses; CW, NF, KS and KL with respect to clinical trial methodology and meta-analysis. Statistical analyses were conducted by EV. The first draft of the manuscript was written by AV and EV. All authors gave comments on early drafts and approved the final version of the manuscript. AV had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. GL died between completion of the first draft of the manuscript and submission to this journal.

Disclosures

An ethics statement was not required for this work. The authors declare that they have no competing interests. The Acupuncture Trialists' Collaboration is funded by an R21 (AT004189I and an R01 (AT006794) from the National Center for Complementary and Alternative Medicine (NCCAM) at the National Institutes of

Health (NIH) to Dr. Vickers) and by a grant from the Samueli Institute. Dr. MacPherson's work on this project was funded in part by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research scheme (RP-PG-0707-10186). Prof. Foster, an NIHR Senior Investigator, was supported through an NIHR Research Professorship (RP-011-015). The views expressed in this publication are those of the author(s) and not necessarily those of the NCCAM, NHS, NIHR or the Department of Health in England. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Accepted Manuscript

Highlights

- Acupuncture has a clinically relevant effect on chronic pain that persists over time
- The effect of acupuncture cannot be explained only by placebo effects
- Factors in addition to the specific effects of needling are important contributors
- Referral for acupuncture treatment is a reasonable option for chronic pain patients

Abstract

Despite wide use in clinical practice, acupuncture remains a controversial treatment for chronic pain. Our objective was to update an individual patient data meta-analysis to determine the effect size of acupuncture for four chronic pain conditions. We searched MEDLINE and the Cochrane Central Registry of Controlled Trials randomized trials published up until December 31, 2015. We included randomized trials of acupuncture needling versus either sham acupuncture or no acupuncture control for non-specific musculoskeletal pain, osteoarthritis, chronic headache, or shoulder pain. Trials were only included if allocation concealment was unambiguously determined to be adequate. Raw data were obtained from study authors and entered into an individual patient data meta-analysis. The main outcome measures were pain and function. An additional 13 trials were identified, with data received for a total of 20,827 patients from 39 trials. Acupuncture was superior to both sham and no acupuncture control for each pain condition (all $p < 0.001$) with differences between groups close to 0.5 standard deviations (SD) for comparison with no acupuncture control and close to 0.2 SDs in comparison with sham. We also found clear evidence that the effects of acupuncture persist over time with only a small decrease, approximately 15%, in treatment effect at one year. In secondary analyses, we found no obvious association between trial outcome and characteristics of acupuncture treatment, but effect sizes of acupuncture were associated with the type of control group, with smaller effects sizes for sham controlled trials that used a penetrating needle for sham, and for trials that had high intensity of intervention in the control arm. We conclude that acupuncture is effective for the treatment of chronic pain, with treatment effects persisting over time. While factors in addition to the specific effects of needling at correct acupuncture point locations are important contributors to the treatment effect, decreases in pain following acupuncture cannot be explained solely in terms of placebo effects. Variations in the effect size of acupuncture in different trials are driven predominately by differences in treatments received by the control group rather than by differences in the characteristics of acupuncture treatment.

Perspective

Acupuncture is effective for the treatment of chronic musculoskeletal, headache and osteoarthritis pain. Treatment effects of acupuncture persist over time and cannot be explained solely in terms of placebo effects. Referral for a course of acupuncture treatment is a reasonable option for a patient with chronic pain.

Keywords: Acupuncture, chronic pain, meta-analysis, osteoarthritis, back pain, neck pain, migraine, tension-type headaches, shoulder pain

Accepted Manuscript

Introduction

Acupuncture remains a controversial treatment for chronic pain, largely due to a provenance outside biomedicine. Traditional acupuncture theory invokes non-anatomical structures such as meridians and non-physiological processes such as the flow of *qi* energy. Although many contemporary practitioners do not rely on such concepts, there remains a dearth of data on how insertion of needles at specific points on the body could lead to long-term decreases in pain. Acupuncture undoubtedly has short-term physiological effects, several of which are relevant to pain^{7, 76, 119}, but there is as yet no explanation as to how such effects could persist.

We previously reported an individual patient data meta-analysis of high-quality trials of acupuncture for chronic pain.⁹² Differences between acupuncture and control in trials without sham (placebo) control were both statistically and clinically significant. Acupuncture was significantly superior to sham control, suggesting that acupuncture effects are not solely explicable in terms of placebo, although these differences were relatively modest. We have separately reported secondary analyses examining whether characteristics of acupuncture treatment⁶⁵ or control groups⁶⁸ influence effect size, and whether the effects of acupuncture treatment persist over time⁶⁹. Here we update our prior analyses now including studies published during the last 7 years.

Methods

The full protocol of the meta-analysis⁹³ and the results of the first individual patient data meta-analysis including RCTs published up to November 2008⁹² have been published. The literature search was repeated to identify eligible RCTs published between December 2008 and December 2015. Trials were considered eligible if they accrued patients with nonspecific back or neck pain, shoulder pain, chronic headache, or osteoarthritis; pain duration was at least 4 weeks for musculoskeletal disorders; at least one group received acupuncture needling and one group received either sham acupuncture or no acupuncture control; the primary endpoint was measured more than 4 weeks after the initial acupuncture treatment; and allocation concealment was determined unambiguously to be adequate. Principal investigators of eligible studies were asked to provide raw data. These raw data were used to replicate all analyses published in the original RCT publication to ensure data accuracy. Each trial was reanalyzed by analysis of covariance with the standardized primary endpoint (scores divided by pooled standard deviation) as the dependent variable, and the baseline measure of the primary endpoint and variables used to stratify randomization as covariates. The primary outcome for each study was that identified by the responding author of each study. The effect sizes for each study were then entered into a meta-analysis using the *metan* command in Stata (version 13, StataCorp, College Station, TX). Both fixed effects and random effects estimates were calculated. Fixed effects weights were calculated using inverse-variance weighting, and random effects weights were calculated using the DerSimonian and Laird method. We pre-specified that meta-analyses would be conducted separately for comparisons of acupuncture vs. sham and acupuncture vs. no acupuncture control, and within each pain type, and the hypothesis test would be based on the fixed effects analysis. In

the original paper, trials for which individual patient data were not available were included as a sensitivity analysis; in this update, we include summary data for these trials in the main meta-analysis and exclude them as a sensitivity analysis.

As secondary analyses, we examined whether characteristics of acupuncture treatment modified treatment effects. Both trial-level and patient-level analyses were performed. For trial-level analyses, we used random-effects meta-regression to test the effect of each characteristic on the main effect estimate using the Stata command *metareg*. For patient-level analyses, we created a linear regression as for the main analysis of effect size, but included the characteristic and an interaction term between the characteristic and treatment allocation. The coefficient was then entered into a meta-analysis. In both analyses, random effects estimates and 95% confidence intervals were reported; p values are based on the fixed effects analysis. We also analyzed the effect of acupuncture relative to different types of sham acupuncture and different types of no acupuncture control group. Three comparisons of sham acupuncture were investigated: penetrating needle vs both non-penetrating needle and non-needle sham; non-penetrating needle vs non-needle sham; and the use of true acupuncture points vs non-acupuncture points among trials using non-penetrating or non-needle sham. For sham arms using penetrating needles, there was also a comparison done between the use of deep needle penetration and shallow needle penetration. We entered the effect size and standard error for each trial into a meta-regression along with the type of sham acupuncture used in that trial. For this analysis, smaller effect sizes indicate a smaller difference in effect between verum acupuncture and sham acupuncture, implying that the type of sham acupuncture used is more active and therefore more similar to verum acupuncture. For the analysis of acupuncture effect relative to no acupuncture control group, we used meta-regression to compare the effects of trials using no acupuncture control groups characterized as high intensity, usual care, or low intensity. We also repeated our prior analyses exploring possible effects of publication bias and exploring difference between sham acupuncture and no treatment.

Results

Systematic Review

Our systematic review⁹³ was updated to include trials published after November 2008 and before December 31, 2015. We identified 75 additional RCTs, of which 13 were eligible (Figure 1). These 13 studies include four trials^{19, 56, 75, 85} included as summary data only in a sensitivity analysis in our first report.

Data Extraction and Quality Assessment

Individual patient data for 2,905 patients were received from 10 of these 13 studies and included patients from the United States, Australia, China, Germany and the UK. For one of the three studies for which we did not receive data, the statisticians involved in the RCT failed to respond to repeated enquiries despite approval for data sharing being obtained from the principal investigator. For the other two studies, the trial authors were contacted and invited to participate but we received no further response. These three studies were included in the analysis as summary data only using the published estimates of effect size.^{31, 70, 75} Two trials from the original systematic review for which data were not received were also included as summary data in these analyses.^{23, 74}

A total of 20,827 patients were included in the total 39 trials (Table 1). The trials comprised 25 comparisons with 16,041 patients of acupuncture and no acupuncture control, and 26 comparisons with 7,237 patients of acupuncture and sham acupuncture control. Of the trials on musculoskeletal pain, most had an eligibility criterion of a minimum 3 or 6 months pain duration. Amongst those for which individual patient data on chronicity were available, the median duration was 4 years (quartiles: 1.1 years, 10 years). There were two trials for which the time period between first symptom and evaluation of outcome could theoretically have been less than three months based on eligibility criteria and timing of assessment. For Irnich et al., the duration of disease was “4 – 52 weeks” for 19% of patients and longer than one year for the remainder.⁴¹ In the case of Kleinhenz et al., no data were provided on chronicity, however, the indication was rotator cuff tendinitis, which is rarely treated in the acute phase.⁵² We conclude that all but a trivial proportion of patients included in the analysis would have met the conventional definition of chronic pain, that is, pain lasting at least 3 to 6 months. Six sham RCTs were determined to have an intermediate likelihood of bias from unblinding.^{13, 26, 41, 49, 59, 103} In one trial, two types of sham acupuncture were used, although only one type (non-needle sham acupuncture) was found to have an intermediate likelihood of bias from unblinding.¹⁰³ One trial (Hinman et al.) was determined to have a sham acupuncture arm with a high likelihood of bias from unblinding.³⁹ This trial was excluded from the main analyses comparing acupuncture to sham acupuncture, but a sensitivity analysis including this trial was performed. None of the 10 new trials included in this analysis had dropout rates of higher than 25%.

Meta-analysis

Forest plots for acupuncture against sham acupuncture and against no acupuncture control are shown separately for each of the 4 pain conditions in Figure 2 and Figure 3. Fixed effects weights are reported in Figures 2 and 3; forest plots with random effects weights reported are presented in Figures S1 and S2 of the supplementary materials. Meta-analytic statistics are shown in Table 2. Consistent with the results of the originally published meta-analysis, acupuncture is found to be statistically superior to control for all analyses ($p < 0.001$). Effect sizes in the updated analyses are similar to those in the original analyses, with effect sizes changing by 0.02 or less for most comparisons. Effect sizes are close to 0.5 in comparison to no acupuncture control and 0.2 for comparisons with sham. To illustrate these effect sizes in more

clinically applicable terms, if baseline pain score in a typical RCT was 60 on a scale of 0-100, with a standard deviation of 25, follow-up scores might be 43 in a no acupuncture control group, 35 in a sham acupuncture group, and 30 among true acupuncture patients. If response were defined as a pain reduction of 50% or more, response rates would be approximately 30%, 42.5% and 50%, respectively. Also in keeping with the original analyses, significant heterogeneity was found in 5 out of 7 comparisons. Significant heterogeneity remained for sham-controlled musculoskeletal pain and osteoarthritis ($p=0.001$ and $p<0.001$, respectively) even after excluding the outlying Vas et al. trials. There was also significant heterogeneity for all indications in the comparison of acupuncture with no acupuncture control. Heterogeneity is further explored below ("Modifiers of Trial Outcome").

Accepted Manuscript

Sensitivity Analyses

Prespecified sensitivity analyses are also shown in Table 2. The exclusion of the RCTs by Vas et al.⁸⁹⁻⁹¹ repeats our prior finding that the effect sizes for comparison with sham are similar for musculoskeletal pain, osteoarthritis and chronic headache. However, there are now sufficient trials for a meta-analysis of shoulder pain trials without inclusion of Vas et al.⁹⁰ and the effect size for this indication is clearly much greater. There is also a large effect size for sham controlled neck pain trials when these are analyzed separately from back pain. Most other sensitivity analyses had little impact on the main findings. Analyses incorporating assessment of patient blinding, missing data or trials without individual patient data, all had very similar results to the primary analysis. As the primary outcome included in the analysis was the outcome specified by the trial authors, we also performed a sensitivity analysis restricted to a single endpoint (pain intensity) at a fixed follow-up time (2 – 3 months after randomization). Results were again very similar apart from sham-controlled trials of musculoskeletal pain (Table 3), where effect size decreased from 0.30 to 0.13, but this appears to be attributable to there being only 5 out of 11 trials that measured pain intensity at 2-3 months, and the trials excluded happened to be those with the larger effect sizes.

We combined all trials into one meta-analysis for all indications to assess the possible effect of publication bias. As in the original analyses, we found some evidence that smaller studies had larger effect sizes for the sham comparison ($p=0.024$), but not for the no acupuncture comparisons ($p=0.75$). No significant asymmetry was seen after excluding the Vas trials and shoulder pain trials from the sham comparison ($N=21$, $p=0.13$), and also when excluding any trials with fewer than 100 patients ($N=21$, $p=0.069$). We found that the difference between acupuncture and control would become non-significant only if there were 51 and >100 unpublished trials with 100 patients and effect sizes in favor of control of 0.25 SD for sham and no acupuncture control respectively.

We also repeated our exploratory analysis comparing sham control with no acupuncture control. In a meta-analysis of 12 RCTs that had both sham and no acupuncture control arms, the effect sizes for sham were 0.39 (95% CI 0.33, 0.45) and 0.45 (95% CI 0.29, 0.61) for fixed and random effects, respectively ($p<0.0001$ for tests of both effect and heterogeneity).

Modifiers of trial outcome

In addition to updating the primary analyses, we also updated previously published analyses on how characteristics of the acupuncture and control interventions influence trial outcomes. Trial-level and patient-level characteristics are found in Tables 4 and 5, respectively.

Acupuncture Characteristics Analysis

We updated previously reported analyses examining whether characteristics of acupuncture treatment modified the effect of acupuncture relative to control. These analyses include both trial-level analysis, based on characteristics described in the study protocol, and patient-level analyses, based on data related to the individual patient. The results are shown in Table 6. We did not find any obvious association between trial outcome and characteristics such as the style of acupuncture (Traditional or Western), use of fixed versus individualized point selection or the use of electrical stimulation. The only clear finding was a dose-response effect to number of acupuncture treatments in trials with a no acupuncture control group (increase in effect size of 0.10 per five sessions, 95% CI -0.01, 0.21, $p=0.001$).

Sham Acupuncture Control Analysis

We also updated a previously published analysis looking at the effects of acupuncture relative to different types of sham acupuncture and no acupuncture control groups. Differences in effect between acupuncture and the different sham acupuncture groups are found in Table 7. The largest difference in effect between acupuncture and sham acupuncture was seen in trials using non-penetrating needles, while the smallest difference was seen in trials using needle penetration. Significant differences were found between trials using penetrating needle sham and those trials that used non-penetrating or non-needle sham (difference in SD -0.30, 95% CI -0.60, -0.00, $p=0.047$), although this result was sensitive to the exclusion of the outlying Vas trials (difference in SD -0.07, 95% CI -0.24, 0.10, $p=0.4$, Table 8), two of which used non-penetrating controls.

No Acupuncture Control Analysis

In addition to updating the analysis comparing types of sham acupuncture control, we also updated the analysis comparing types of no acupuncture control. We updated the categorization of no acupuncture control groups, and categorized trials as having a high intensity, usual care, or low intensity control group. In a “high intensity” control group, patients received a specified course of protocol-guided treatment. For instance, the UK APEX trial by Foster et al.³³ is considered a high intensity control because patients were randomized to receive a course of individualized, supervised physical therapy plus acupuncture vs. physical therapy alone. In a trial with “usual care” control, patients are able to access whatever care they might reasonably receive outside of the study. As an example, in the UK NHS study, patients were randomized to “use” vs. “avoid” acupuncture and could receive whatever other treatments were offered to them.⁹⁵ A control group was defined as “low intensity” if patients were not allowed to receive certain treatments that might otherwise be available. For instance, the Acupuncture Randomized Trials for low back pain and osteoarthritis limited treatment of pain in the control group to oral nonsteroidal anti-inflammatory drugs, excluding other types of treatment, such as steroids and other classes of analgesics.^{11, 108} Trials were assessed and assigned a control group type by three collaborators, with disagreements resolved by consensus. One trial was excluded from this analysis as there was a reasonable argument that it involved active control, prespecified to be excluded.²⁶ Differences in effect between acupuncture and no acupuncture control groups are presented in Table 7. Significant differences were found between

acupuncture and control for all types of no acupuncture control group. Notably, however, in trials that had high intensity control groups, acupuncture had smaller effect sizes compared to those with low intensity controls groups (difference - 0.81, 95% CI -1.26, -0.36, $p=0.0004$); similarly in trials with usual care control acupuncture had smaller effect sizes than trials with a low intensity control group (difference in SD -0.65, 95% CI -0.98, -0.31, $p=0.0002$, Table 8).

Time Course of Acupuncture Effects Analysis

We updated a previously published analysis assessing change in the effects of acupuncture over time relative to sham acupuncture and no acupuncture control⁶⁹. Number of weeks of acupuncture treatment and the time points used in this analysis are reported in Table 9. A total of 14 trials and 4,124 patients were included in the analysis of acupuncture vs no acupuncture control. The fixed-effects estimate for the between-group comparison of acupuncture vs no acupuncture controls showed a decrease in the effect size of acupuncture of 0.019 SD per 3 months (95% CI -0.041, 0.003, $p=0.096$, $p=0.011$ for heterogeneity, Figure 4a). Given a difference between acupuncture and no acupuncture control of around 0.5 SD, this is equivalent to about a 15% decrease in acupuncture effect relative to control at 1 year after randomization, which was usually between 9 and 10 months after the end of treatment. In the analysis of acupuncture vs sham acupuncture, a total of 21 trials and 6,276 patients were included. There was a non-significant decrease of 0.012 SD per 3 months in acupuncture relative to sham acupuncture (95% CI -0.035, 0.011, $p=0.3$, Figure 4b), about a 25% decrease in acupuncture effect at 1 year after randomization. Significant heterogeneity among trials was seen ($p<0.0001$). The previous analysis found that the decrease in effect of acupuncture relative to sham was driven by the decrease in neck pain trials (a decrease of 0.587 SD per 3 months, 95% CI -0.767, -0.406, $p<0.0001$). We also analyzed the change in acupuncture relative to sham excluding these trials and found a non-significant decrease of -0.003 SD per 3 months (95% CI -0.026, 0.020, $p=0.8$) with no significant heterogeneity among trials ($p=0.12$). Hence almost all the decrease in acupuncture effects in this analysis seems attributable to neck pain.

As a sensitivity analysis, we repeated the analyses including only trials that found a significant difference between acupuncture and control, as trials that showed no difference between groups cannot show a reduction in acupuncture effects over time. Nine trials with 2,997 patients were included in this analysis for the comparison between acupuncture and no acupuncture controls. A smaller and still non-significant decrease in the effect of acupuncture was found (-0.008 SD per 3 months, 95% CI -0.034, 0.018, $p=0.5$) and heterogeneity between trials was reduced ($p=0.082$). None of the newly included trials showed a significant effect of acupuncture vs sham and so this analysis of sham-controlled trials with a significant effect contains the same 7 trials and 1,450 patients and has the same results as reported in the original publication (-0.049 SD per 3 months, 95% CI -0.086, -0.013, $p=0.008$, heterogeneity $p<0.0001$).

Discussion

We updated an individual patient data meta-analysis of high-quality trials of acupuncture for chronic pain with seven additional years of data. An additional 10 studies were included with nearly 3,000 patients. In total, our analyses include 39 studies and 20,827 patients. The results confirm and strengthen prior key findings that acupuncture has a clinically relevant effect compared to no acupuncture control. Moreover, we confirmed that, although the effects of acupuncture are not completely explicable in terms of placebo effects, factors other than the specific effects of needling at correct acupuncture point locations are important contributors to acupuncture treatment benefit. Effects of acupuncture appear to persist over at least a 12 month period.

Heterogeneity continues to be an obvious aspect of our findings, with the results of trials varying by more than would be expected by chance. We have presented data that heterogeneity is predominately driven by differences between control groups rather than by differences between acupuncture treatment characteristics. We did not find any obvious differences between the results of trials depending on treatment characteristics such as style of acupuncture, duration of treatment sessions or training of acupuncturists. By contrast, we found evidence that effect sizes of acupuncture were smaller for sham-controlled trials with penetrating needles and for no acupuncture control trials where patients received high intensity care (e.g. a trial of acupuncture plus physical therapy vs. physical therapy alone). In some cases, heterogeneity was also driven by a set of outlying trials with large effect sizes. We have presented these analyses with and without the outlying trials to provide all necessary information for interpreting these results and drawing conclusions.

Another novel finding is the higher than average effects of acupuncture on upper body musculoskeletal pain. We now have sufficient data to conduct a meta-analysis for neck pain and for shoulder pain, even after exclusion of outlying trials. The effect sizes versus sham, 0.57 for shoulder and 0.83 for neck pain, were much larger than seen for low back pain, osteoarthritis and headache, although we also saw evidence that treatment benefits did not persist for neck pain.

Since publication of our results, there has been no substantive critique of our methodology in the peer-reviewed literature. The main issue under discussion seems to be whether the effect size of acupuncture is clinically relevant⁹⁴, specifically, whether clinical relevance is determined by the comparison with no acupuncture control or by comparison with sham. We have previously argued in favor of the former, on the grounds that the clinical decision made by a referring clinician in discussion with their patient is not between acupuncture and sham but between acupuncture and no acupuncture. Our argument is given the context of the excellent safety profile of acupuncture⁶⁶, evidence that the non-specific effects of acupuncture are particular to acupuncture and are not easily reproduced^{46, 54} and evidence provided here and elsewhere⁹ that some interventions used as sham acupuncture may be physiologically active.

It is also illustrative to compare our results to those of other interventions routinely used in clinical practice. For instance, in one meta-analysis of non-steroidal anti-inflammatory drugs (NSAIDs) for osteoarthritis of the knee, the

effect size for NSAIDs vs placebo for trials that did not preselect NSAID responders was 0.23;¹⁰ for chronic low back pain, the effect size for NSAIDs is < 0.20 ²⁹.

We find several implications for research. In terms of the methodology of subsequent acupuncture trials for chronic pain, we find that the balance of evidence is to give a higher dose of acupuncture in terms of a greater number of treatments in trials without sham control. Although the nature of the control group in trials will naturally be driven by the research question, investigators should be aware of the evidence that control arms that incorporate a relatively intense level of intervention, such as when acupuncture is added into an intensive rehabilitation regimen, tend to lead to smaller effect sizes, as do sham controls that involve needle penetration. Further research is warranted on whether acupuncture is particularly effective for upper body musculoskeletal pain. An associated hypothesis is whether there are subtypes of other chronic pain indications that have differential response to acupuncture. It would naturally be ideal to know before referring a patient for treatment whether, say, the type of back pain they are experiencing is one that would be amenable to treatment with acupuncture. We will also repeat our prior call for research on how best to incorporate acupuncture into the multidisciplinary care of chronic pain patients.

In terms of implications for clinical practice, we have confirmed that acupuncture has a clinically relevant, persistent effect on chronic pain that is not completely explained by placebo effects. Referral for a course of acupuncture treatment is therefore a reasonable option for a patient with chronic pain.

Acupuncture Trialists' Collaboration Members

Claire Allen, BA, Evidence Aid, Oxford, United Kingdom.

Brian Berman, MD, University of Maryland School of Medicine and Center for Integrative Medicine, College Park, Maryland.

Benno Brinkhaus, MD, Institute for Social Medicine, Epidemiology and Health Economics, Charité University Medical Center, Berlin, Germany.

Remy Coeytaux, MD, PhD, Department of Community and Family Medicine, Duke University, Durham, North Carolina.

Hans-Christoph Diener, MD, PhD, Department of Neurology, University of Duisburg-Essen, Germany.

Heinz G. Endres, MD, Ruhr-University Bochum, Bochum, Germany.

Nadine E. Foster, DPhil, BSc(Hons), Arthritis Research UK Primary Care Centre, Research Institute of Primary Care and Health Sciences, Keele University, Newcastle-under-Lyme, Staffordshire, England.

Michael Haake, MD, PhD, Department of Orthopedics and Traumatology, SLK Hospitals, Heilbronn, Germany.

Rana S. Hinman, PhD, University of Melbourne, Melbourne, Australia.

Dominik Irnich, MD, Multidisciplinary Pain Centre, Department of Anesthesiology, Ludwig-Maximilians-Universität München (LMU Munich), Germany.

Wayne B. Jonas, MD, Samueli Institute, Alexandria, Virginia.

Kai Kronfeld, PhD, Interdisciplinary Centre for Clinical Trials (IZKS Mainz), University Medical Centre Mainz, Mainz, Germany.

Lixing Lao, PhD, University of Maryland and Center for Integrative Medicine, College Park, Maryland.

George Lewith, MD, FRCP, Primary Care and Population Sciences, Faculty of Medicine, University of Southampton, Southampton, England.

Klaus Linde, MD, Institute of General Practice, Technical University Munich, Munich, Germany.

Hugh MacPherson, PhD, Professor of Acupuncture Research, Department of Health Sciences, University of York, York, England.

Eric Manheimer, MS, Center for Integrative Medicine, University of Maryland School of Medicine, College Park, Maryland.

Dieter Melchart, MD, PhD, Competence Centre for Complementary Medicine and Naturopathy, Technical University Munich, Munich, Germany.

Albrecht Molsberger, MD, PhD, German Acupuncture Research Group, Duesseldorf, Germany.

Karen J. Sherman, PhD, MPH, Group Health Research Institute, Seattle, Washington.

Maria Suarez-Almazor, MD, PhD, MD Anderson Cancer Center, Houston, Texas.

Hans Trampisch, PhD, Department of Medical Statistics and Epidemiology, Ruhr–University Bochum, Germany.

Jorge Vas, MD, PhD, Pain Treatment Unit, Dos Hermanas Primary Care Health Center (Andalusia Public Health System), Dos Hermanas, Spain.

Andrew J. Vickers (collaboration chair), DPhil, Memorial Sloan Kettering Cancer Center, New York, New York

Peter White, PhD, School of Health Sciences, University of Southampton, England.

Lyn Williamson, MD, MA (Oxon), MRCP, FRCP, Great Western Hospital, Swindon, UK.

Stefan N. Willich, MD, MPH, MBA, Institute for Social Medicine, Epidemiology and Health Economics, Charité University Medical Center, Berlin, Germany.

Claudia M. Witt, MD, MBA, Institute for Complementary and Integrative Medicine, University of Zurich and University Hospital Zurich, Zurich, Switzerland; Institute for Social Medicine, Epidemiology and Health Economics, Charite-Universitätsmedizin, Berlin, Germany; Center for Integrative Medicine, University of Maryland School of Medicine, Baltimore, Maryland.

References

1. Ahn CB, Lee SJ, Lee JC, Fossion JP, Sant'Ana A. A clinical pilot study comparing traditional acupuncture to combined acupuncture for treating headache, trigeminal neuralgia and retro-auricular pain in facial palsy. *Journal of acupuncture and meridian studies*. 4:29-43, 2011
2. Ahsin S, Saleem S, Bhatti AM, Iles RK, Aslam M. Clinical and endocrinological changes after electro-acupuncture treatment in patients with osteoarthritis of the knee. *Pain*. 147:60-66, 2009
3. Allais G, Romoli M, Rolando S, Airola G, Castagnoli Gabellari I, Allais R, Benedetto C. Ear acupuncture in the treatment of migraine attacks: a randomized trial on the efficacy of appropriate versus inappropriate acupoints. *Neurological sciences : official journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology*. 32 Suppl 1:S173-175, 2011
4. Allam H, Mohammed NH. The Role of Scalp Acupuncture for Relieving the Chronic Pain of Degenerative Osteoarthritis: A Pilot Study of Egyptian Women. *Medical acupuncture*. 25:216-220, 2013
5. Aranha MF, Muller CE, Gaviao MB. Pain intensity and cervical range of motion in women with myofascial pain treated with acupuncture and electroacupuncture: a double-blinded, randomized clinical trial. *Brazilian journal of physical therapy*. 0:0, 2014
6. Ashraf A, Zarei F, Hadianfard MJ, Kazemi B, Mohammadi S, Naseri M, Nasser A, Khodadadi M, Sayadi M. Comparison the effect of lateral wedge insole and acupuncture in medial compartment knee osteoarthritis: a randomized controlled trial. *The Knee*. 21:439-444, 2014
7. Baeumler PI, Fleckenstein J, Takayama S, Simang M, Seki T, Irnich D. Effects of acupuncture on sensory perception: a systematic review and meta-analysis. *PloS one*. 9:e113731, 2014
8. Berman BM, Lao L, Langenberg P, Lee WL, Gilpin AM, Hochberg MC. Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: a randomized, controlled trial. *Ann Intern Med*. 141:901-910, 2004
9. Birch S. A review and analysis of placebo treatments, placebo effects, and placebo controls in trials of medical procedures when sham is not inert. *J Altern Complement Med*. 12:303-310, 2006
10. Bjordal JM, Ljunggren AE, Klovning A, Slordal L. Non-steroidal anti-inflammatory drugs, including cyclo-oxygenase-2 inhibitors, in osteoarthritic knee pain: meta-analysis of randomised placebo controlled trials. *BMJ*. 329:1317, 2004
11. Brinkhaus B, Witt CM, Jena S, Linde K, Streng A, Wagenpfeil S, Irnich D, Walther HU, Melchart D, Willich SN. Acupuncture in patients with chronic low back pain: a randomized controlled trial. *Arch Intern Med*. 166:450-457, 2006
12. Cameron ID, Wang E, Sindhusake D. A randomized trial comparing acupuncture and simulated acupuncture for subacute and chronic whiplash. *Spine (Phila Pa 1976)*. 36:E1659-1665, 2011
13. Carlsson CP, Sjolund BH. Acupuncture for chronic low back pain: a randomized placebo-controlled study with long-term follow-up. *Clin J Pain*. 17:296-305, 2001
14. Chassot M, Dussan-Sarria JA, Sehn FC, Deitos A, de Souza A, Vercelino R, Torres IL, Fregni F, Caumo W. Electroacupuncture analgesia is associated with increased serum brain-derived neurotrophic factor in chronic tension-type headache: a randomized, sham controlled, crossover trial. *BMC complementary and alternative medicine*. 15:144, 2015
15. Chen CC, Yang CC, Hu CC, Shih HN, Chang YH, Hsieh PH. Acupuncture for pain relief after total knee arthroplasty: a randomized controlled trial. *Regional anesthesia and pain medicine*. 40:31-36, 2015
16. Chen LX, Mao JJ, Fernandes S, Galantino ML, Guo W, Lariccia P, Teal VL, Bowman MA, Schumacher HR, Farrar JT. Integrating acupuncture with exercise-based physical therapy for knee osteoarthritis: a randomized controlled trial. *Journal of clinical rheumatology : practical reports on rheumatic & musculoskeletal diseases*. 19:308-316, 2013
17. Chen X, Spaeth RB, Freeman SG, Scarborough DM, Hashmi JA, Wey HY, Egorova N, Vangel M, Mao J, Wasan AD, Edwards RR, Gollub RL, Kong J. The modulation effect of longitudinal acupuncture on resting state functional connectivity in knee osteoarthritis patients. *Molecular pain*. 11:67, 2015
18. Cherkin DC, Eisenberg D, Sherman KJ, Barlow W, Kaptchuk TJ, Street J, Deyo RA. Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and self-care education for chronic low back pain. *Arch Intern Med*. 161:1081-1088, 2001

19. Cherkin DC, Sherman KJ, Avins AL, Erro JH, Ichikawa L, Barlow WE, Delaney K, Hawkes R, Hamilton L, Pressman A, Khalsa PS, Deyo RA. A randomized trial comparing acupuncture, simulated acupuncture, and usual care for chronic low back pain. *Arch Intern Med.* 169:858-866, 2009
20. Cho JH, Nam DH, Kim KT, Lee JH. Acupuncture with non-steroidal anti-inflammatory drugs (NSAIDs) versus acupuncture or NSAIDs alone for the treatment of chronic neck pain: an assessor-blinded randomised controlled pilot study. *Acupunct Med.* 32:17-23, 2014
21. Cho YJ, Song YK, Cha YY, Shin BC, Shin IH, Park HJ, Lee HS, Kim KW, Cho JH, Chung WS, Lee JH, Song MY. Acupuncture for chronic low back pain: a multicenter, randomized, patient-assessor blind, sham-controlled clinical trial. *Spine (Phila Pa 1976).* 38:549-557, 2013
22. Chou LW, Hsieh YL, Chen HS, Hong CZ, Kao MJ, Han TI. Remote therapeutic effectiveness of acupuncture in treating myofascial trigger point of the upper trapezius muscle. *American journal of physical medicine & rehabilitation / Association of Academic Physiatrists.* 90:1036-1049, 2011
23. Coeytaux RR, Kaufman JS, Kaptchuk TJ, Chen W, Miller WC, Callahan LF, Mann JD. A randomized, controlled trial of acupuncture for chronic daily headache. *Headache.* 45:1113-1123, 2005
24. D'Alessandro E, de Brito C, Cecatto R, Saul M, Atta JA, Lin CA. Evaluation of acupuncture for cancer symptoms in a cancer institute in Brazil. *Acupunct Med.* 31:23-26, 2013
25. De Kooning M, Tobbackx Y, Meeus M, Wauters L, Ickmans K, De Vilder P, Roose J, Verhaeghe T, Nijs J. Acupuncture-Analgesia Following a Single Treatment Session in Chronic Whiplash is Unrelated to Autonomic Nervous System Changes: A Randomized Cross-over Trial. *Pain physician.* 18:527-536, 2015
26. Diener HC, Kronfeld K, Boewing G, Lungenhausen M, Maier C, Molsberger A, Tegenthoff M, Trampisch HJ, Zenz M, Meinert R. Efficacy of acupuncture for the prophylaxis of migraine: a multicentre randomised controlled clinical trial. *Lancet Neurol.* 5:310-316, 2006
27. Egorova N, Gollub RL, Kong J. Repeated verum but not placebo acupuncture normalizes connectivity in brain regions dysregulated in chronic pain. *NeuroImage. Clinical.* 9:430-435, 2015
28. Endres HG, Bowing G, Diener HC, Lange S, Maier C, Molsberger A, Zenz M, Vickers AJ, Tegenthoff M. Acupuncture for tension-type headache: a multicentre, sham-controlled, patient-and observer-blinded, randomised trial. *J Headache Pain.* 8:306-314, 2007
29. Enthoven WT, Roelofs PD, Deyo RA, van Tulder MW, Koes BW. Non-steroidal anti-inflammatory drugs for chronic low back pain. *Cochrane Database Syst Rev.* 2:CD012087, 2016
30. Facco E, Liguori A, Petti F, Fauci AJ, Cavallin F, Zanette G. Acupuncture versus valproic acid in the prophylaxis of migraine without aura: a prospective controlled study. *Minerva anesthesiologica.* 79:634-642, 2013
31. Ferro EC, Biagini AP, da Silva IE, Silva ML, Silva JR. The combined effect of acupuncture and Tanacetum parthenium on quality of life in women with headache: randomised study. *Acupunct Med.* 30:252-257, 2012
32. Foroughipour M, Golchian AR, Kalhor M, Akhlaghi S, Farzadfard MT, Azizi H. A sham-controlled trial of acupuncture as an adjunct in migraine prophylaxis. *Acupunct Med.* 32:12-16, 2014
33. Foster NE, Thomas E, Barlas P, Hill JC, Young J, Mason E, Hay EM. Acupuncture as an adjunct to exercise based physiotherapy for osteoarthritis of the knee: randomised controlled trial. *Bmj.* 335:436, 2007
34. Grotle M. Traditional Chinese acupuncture was not superior to sham acupuncture for knee osteoarthritis but delivering treatment with high expectations of improvement was superior to delivering treatment with neutral expectations. *Journal of physiotherapy.* 57:56, 2011
35. Guerra de Hoyos JA, Andres Martin Mdel C, Bassas y Baena de Leon E, Vigara Lopez M, Molina Lopez T, Verdugo Morilla FA, Gonzalez Moreno MJ. Randomised trial of long term effect of acupuncture for shoulder pain. *Pain.* 112:289-298, 2004
36. Haake M, Muller HH, Schade-Brittinger C, Basler HD, Schafer H, Maier C, Endres HG, Trampisch HJ, Molsberger A. German Acupuncture Trials (GERAC) for chronic low back pain: randomized, multicenter, blinded, parallel-group trial with 3 groups. *Arch Intern Med.* 167:1892-1898, 2007
37. Hansson Y, Carlsson C, Olsson E. Intramuscular and periosteal acupuncture in patients suffering from chronic musculoskeletal pain - a controlled trial. *Acupunct Med.* 26:214-223, 2008
38. Hasegawa TM, Baptista AS, de Souza MC, Yoshizumi AM, Natour J. Acupuncture for acute non-specific low back pain: a randomised, controlled, double-blind, placebo trial. *Acupunct Med.* 32:109-115, 2014

39. Hinman RS, McCrory P, Pirotta M, Relf I, Forbes A, Crossley KM, Williamson E, Kyriakides M, Novy K, Metcalf BR, Harris A, Reddy P, Conaghan PG, Bennell KL. Acupuncture for chronic knee pain: a randomized clinical trial. *Jama*. 312:1313-1322, 2014
40. Hunter RF, McDonough SM, Bradbury I, Liddle SD, Walsh DM, Dhamija S, Glasgow P, Gormley G, McCann SM, Park J, Hurley DA, Delitto A, Baxter GD. Exercise and Auricular Acupuncture for Chronic Low-back Pain: A Feasibility Randomized-controlled Trial. *Clin J Pain*. 28:259-267, 2012
41. Irnich D, Behrens N, Molzen H, Konig A, Gleditsch J, Krauss M, Natalis M, Senn E, Beyer A, Schops P. Randomised trial of acupuncture compared with conventional massage and "sham" laser acupuncture for treatment of chronic neck pain. *Bmj*. 322:1574-1578, 2001
42. Itoh K, Saito S, Sahara S, Naitoh Y, Imai K, Kitakoji H. Randomized trial of trigger point acupuncture treatment for chronic shoulder pain: a preliminary study. *Journal of acupuncture and meridian studies*. 7:59-64, 2014
43. Jena S, Witt CM, Brinkhaus B, Wegscheider K, Willich SN. Acupuncture in patients with headache. *Cephalalgia*. 28:969-979, 2008
44. Ji L, Wang H, Cao Y, Yan P, Jin X, Nie P, Wang C, Li R, Zhang C, Yang M, Yang J. Sharp-Hook Acupuncture (Feng Gou Zhen) for Patients with Periarthritis of Shoulder: A Randomized Controlled Trial. *Evid Based Complement Alternat Med*. 2015:312309, 2015
45. Johansson K, Bergstrom A, Schroder K, Foldevi M. Subacromial corticosteroid injection or acupuncture with home exercises when treating patients with subacromial impingement in primary care--a randomized clinical trial. *Family practice*. 28:355-365, 2011
46. Kaptchuk TJ, Stason WB, Davis RB, Legedza AR, Schnyer RN, Kerr CE, Stone DA, Nam BH, Kirsch I, Goldman RH. Sham device v inert pill: randomised controlled trial of two placebo treatments. *BMJ*. 332:391-397, 2006
47. Karner M, Brazkiewicz F, Remppis A, Fischer J, Gerlach O, Stremmel W, Subramanian SV, Greten HJ. Objectifying specific and nonspecific effects of acupuncture: a double-blinded randomised trial in osteoarthritis of the knee. *Evid Based Complement Alternat Med*. 2013:427265, 2013
48. Kennedy S, Baxter GD, Kerr DP, Bradbury I, Park J, McDonough SM. Acupuncture for acute non-specific low back pain: a pilot randomised non-penetrating sham controlled trial. *Complement Ther Med*. 16:139-146, 2008
49. Kerr DP, Walsh DM, Baxter D. Acupuncture in the management of chronic low back pain: a blinded randomized controlled trial. *Clin J Pain*. 19:364-370, 2003
50. Kim EJ, Lim CY, Lee EY, Lee SD, Kim KS. Comparing the effects of individualized, standard, sham and no acupuncture in the treatment of knee osteoarthritis: a multicenter randomized controlled trial. *Trials*. 14:129, 2013
51. Kim JI, Kim TH, Lee MS, Kang JW, Kim KH, Choi JY, Kang KW, Kim AR, Shin MS, Jung SY, Choi SM. Evaluation of wet-cupping therapy for persistent non-specific low back pain: a randomised, waiting-list controlled, open-label, parallel-group pilot trial. *Trials*. 12:146, 2011
52. Kleinhenz J, Streitberger K, Windeler J, Gussbacher A, Mavridis G, Martin E. Randomised clinical trial comparing the effects of acupuncture and a newly designed placebo needle in rotator cuff tendinitis. *Pain*. 83:235-241, 1999
53. Koh PS, Seo BK, Cho NS, Park HS, Park DS, Baek YH. Clinical effectiveness of bee venom acupuncture and physiotherapy in the treatment of adhesive capsulitis: a randomized controlled trial. *Journal of shoulder and elbow surgery / American Shoulder and Elbow Surgeons ... [et al.]*. 22:1053-1062, 2013
54. Kong J, Spaeth R, Cook A, Kirsch I, Claggett B, Vangel M, Gollub RL, Smoller JW, Kaptchuk TJ. Are all placebo effects equal? Placebo pills, sham acupuncture, cue conditioning and their association. *PLoS one*. 8:e67485, 2013
55. Lan L, Gao Y, Zeng F, Qin W, Dong M, Liu M, Guo T, Liang F. A central analgesic mechanism of acupuncture for migraine: An ongoing functional MRI study. *Neural regeneration research*. 8:2649-2655, 2013
56. Lansdown H, Howard K, Brealey S, MacPherson H. Acupuncture for pain and osteoarthritis of the knee: a pilot study for an open parallel-arm randomised controlled trial. *BMC Musculoskelet Disord*. 10:130, 2009
57. Lathia AT, Jung SM, Chen LX. Efficacy of acupuncture as a treatment for chronic shoulder pain. *J Altern Complement Med*. 15:613-618, 2009
58. Li N, Tian F, Wang C, Yu P, Zhou X, Wen Q, Qiao X, Huang L. Therapeutic effect of acupuncture and massage for shoulder-hand syndrome in hemiplegia patients: a clinical two-center randomized controlled trial. *Journal of*

traditional Chinese medicine = *Chung i tsa chih ying wen pan* / sponsored by All-China Association of Traditional Chinese Medicine, Academy of Traditional Chinese Medicine. 32:343-349, 2012

59. Li Y, Zheng H, Witt CM, Roll S, Yu SG, Yan J, Sun GJ, Zhao L, Huang WJ, Chang XR, Zhang HX, Wang DJ, Lan L, Zou R, Liang FR. Acupuncture for migraine prophylaxis: a randomized controlled trial. *Cmaj*. 184:401-410, 2012
60. Liang Z, Zhu X, Yang X, Fu W, Lu A. Assessment of a traditional acupuncture therapy for chronic neck pain: a pilot randomised controlled study. *Complement Ther Med*. 19 Suppl 1:S26-32, 2011
61. Lin ML, Lin MH, Fen JJ, Lin WT, Lin CW, Chen PQ. A comparison between pulsed radiofrequency and electro-acupuncture for relieving pain in patients with chronic low back pain. *Acupuncture & electro-therapeutics research*. 35:133-146, 2010
62. Lin ML, Wu HC, Hsieh YH, Su CT, Shih YS, Lin CW, Wu JH. Evaluation of the effect of laser acupuncture and cupping with ryodoraku and visual analog scale on low back pain. *Evid Based Complement Alternat Med*. 2012:521612, 2012
63. Linde K, Streng A, Jurgens S, Hoppe A, Brinkhaus B, Witt C, Wagenpfeil S, Pfaffenrath V, Hammes MG, Weidenhammer W, Willich SN, Melchart D. Acupuncture for patients with migraine: a randomized controlled trial. *Jama*. 293:2118-2125, 2005
64. Ma C, Wu S, Li G, Xiao X, Mai M, Yan T. Comparison of miniscalpel-needle release, acupuncture needling, and stretching exercise to trigger point in myofascial pain syndrome. *Clin J Pain*. 26:251-257, 2010
65. MacPherson H, Maschino AC, Lewith G, Foster NE, Witt CM, Vickers AJ. Characteristics of acupuncture treatment associated with outcome: an individual patient meta-analysis of 17,922 patients with chronic pain in randomised controlled trials. *PLoS one*. 8:e77438, 2013
66. Macpherson H, Scullion A, Thomas KJ, Walters S. Patient reports of adverse events associated with acupuncture treatment: a prospective national survey. *Quality & safety in health care*. 13:349-355, 2004
67. MacPherson H, Tilbrook H, Richmond S, Woodman J, Ballard K, Atkin K, Bland M, Eldred J, Essex H, Hewitt C, Hopton A, Keding A, Lansdown H, Parrott S, Torgerson D, Wenham A, Watt I. Alexander Technique Lessons or Acupuncture Sessions for Persons With Chronic Neck Pain: A Randomized Trial. *Ann Intern Med*. 163:653-662, 2015
68. MacPherson H, Vertosick E, Lewith G, Linde K, Sherman KJ, Witt CM, Vickers AJ. Influence of control group on effect size in trials of acupuncture for chronic pain: a secondary analysis of an individual patient data meta-analysis. *PLoS one*. 9:e93739, 2014
69. MacPherson H, Vertosick EA, Foster NE, Lewith G, Linde K, Sherman KJ, Witt CM, Vickers AJ, Acupuncture Trialists C. The persistence of the effects of acupuncture after a course of treatment: A meta-analysis of patients with chronic pain. *Pain*. 2016
70. Mavrommatis CI, Argyra E, Vadalouka A, Vasilakos DG. Acupuncture as an adjunctive therapy to pharmacological treatment in patients with chronic pain due to osteoarthritis of the knee: a 3-armed, randomized, placebo-controlled trial. *Pain*. 153:1720-1726, 2012
71. Melchart D, Streng A, Hoppe A, Brinkhaus B, Witt C, Wagenpfeil S, Pfaffenrath V, Hammes M, Hummelsberger J, Irnich D, Weidenhammer W, Willich SN, Linde K. Acupuncture in patients with tension-type headache: randomised controlled trial. *Bmj*. 331:376-382, 2005
72. Miller E, Maimon Y, Rosenblatt Y, Mendler A, Hasner A, Barad A, Amir H, Dekel S, Lev-Ari S. Delayed Effect of Acupuncture Treatment in OA of the Knee: A Blinded, Randomized, Controlled Trial. *Evid Based Complement Alternat Med*. 2011:792975, 2011
73. Miyazaki S, Hagihara A, Kanda R, Mukaino Y, Nobutomo K. Applicability of press needles to a double-blind trial: a randomized, double-blind, placebo-controlled trial. *Clin J Pain*. 25:438-444, 2009
74. Molsberger AF, Mau J, Pawelec DB, Winkler J. Does acupuncture improve the orthopedic management of chronic low back pain--a randomized, blinded, controlled trial with 3 months follow up. *Pain*. 99:579-587, 2002
75. Molsberger AF, Schneider T, Gotthardt H, Drabik A. German Randomized Acupuncture Trial for chronic shoulder pain (GRASP) - a pragmatic, controlled, patient-blinded, multi-centre trial in an outpatient care environment. *Pain*. 151:146-154, 2010
76. Peets JM, Pomeranz B. CXBK mice deficient in opiate receptors show poor electroacupuncture analgesia. *Nature*. 273:675-676, 1978

77. Sahin N, Ozcan E, Sezen K, Karatas O, Issever H. Efficacy of acupuncture in patients with chronic neck pain--a randomised, sham controlled trial. *Acupuncture & electro-therapeutics research*. 35:17-27, 2010
78. Saleki M, Ahadi T, Razi M, Raeisi GR, Forough B, Ali MK. Comparison of the effects of acupuncture and isometric exercises on symptom of knee osteoarthritis. *International journal of preventive medicine*. 4:S73-77, 2013
79. Salter GC, Roman M, Bland MJ, MacPherson H. Acupuncture for chronic neck pain: a pilot for a randomised controlled trial. *BMC Musculoskelet Disord*. 7:99, 2006
80. Scharf HP, Mansmann U, Streitberger K, Witte S, Kramer J, Maier C, Trampisch HJ, Victor N. Acupuncture and knee osteoarthritis: a three-armed randomized trial. *Ann Intern Med*. 145:12-20, 2006
81. Shankar N, Thakur M, Tandon OP, Saxena AK, Arora S, Bhattacharya N. Autonomic status and pain profile in patients of chronic low back pain and following electro acupuncture therapy: a randomized control trial. *Indian journal of physiology and pharmacology*. 55:25-36, 2011
82. Silva AC, Biasotto-Gonzalez DA, Dos Santos DM, Melo NC, Gomes CA, Amorim CF, Politti F. Evaluation of the Immediate Effect of Auricular Acupuncture on Pain and Electromyographic Activity of the Upper Trapezius Muscle in Patients with Nonspecific Neck Pain: A Randomized, Single-Blinded, Sham-Controlled, Crossover Study. *Evid Based Complement Alternat Med*. 2015:523851, 2015
83. Soderberg EI, Carlsson JY, Stener-Victorin E, Dahlof C. Subjective well-being in patients with chronic tension-type headache: effect of acupuncture, physical training, and relaxation training. *Clin J Pain*. 27:448-456, 2011
84. Spaeth RB, Camhi S, Hashmi JA, Vangel M, Wasan AD, Edwards RR, Gollub RL, Kong J. A longitudinal study of the reliability of acupuncture deqi sensations in knee osteoarthritis. *Evid Based Complement Alternat Med*. 2013:204259, 2013
85. Suarez-Almazor ME, Looney C, Liu Y, Cox V, Pietz K, Marcus DM, Street RL, Jr. A randomized controlled trial of acupuncture for osteoarthritis of the knee: effects of patient-provider communication. *Arthritis Care Res (Hoboken)*. 62:1229-1236, 2010
86. Sun MY, Hsieh CL, Cheng YY, Hung HC, Li TC, Yen SM, Huang IS. The therapeutic effects of acupuncture on patients with chronic neck myofascial pain syndrome: a single-blind randomized controlled trial. *Am J Chin Med*. 38:849-859, 2010
87. Thomas KJ, MacPherson H, Thorpe L, Brazier J, Fitter M, Campbell MJ, Roman M, Walters SJ, Nicholl J. Randomised controlled trial of a short course of traditional acupuncture compared with usual care for persistent non-specific low back pain. *Bmj*. 333:623, 2006
88. Vas J, Aranda JM, Modesto M, Benitez-Parejo N, Herrera A, Martinez-Barquin DM, Aguilar I, Sanchez-Araujo M, Rivas-Ruiz F. Acupuncture in patients with acute low back pain: a multicentre randomised controlled clinical trial. *Pain*. 153:1883-1889, 2012
89. Vas J, Mendez C, Perea-Milla E, Vega E, Panadero MD, Leon JM, Borge MA, Gaspar O, Sanchez-Rodriguez F, Aguilar I, Jurado R. Acupuncture as a complementary therapy to the pharmacological treatment of osteoarthritis of the knee: randomised controlled trial. *Bmj*. 329:1216, 2004
90. Vas J, Ortega C, Olmo V, Perez-Fernandez F, Hernandez L, Medina I, Seminario JM, Herrera A, Luna F, Perea-Milla E, Mendez C, Madrazo F, Jimenez C, Ruiz MA, Aguilar I. Single-point acupuncture and physiotherapy for the treatment of painful shoulder: a multicentre randomized controlled trial. *Rheumatology (Oxford)*. 47:887-893, 2008
91. Vas J, Perea-Milla E, Mendez C, Sanchez Navarro C, Leon Rubio JM, Brioso M, Garcia Obrero I. Efficacy and safety of acupuncture for chronic uncomplicated neck pain: a randomised controlled study. *Pain*. 126:245-255, 2006
92. Vickers AJ, Cronin AM, Maschino AC, Lewith G, MacPherson H, Foster NE, Sherman KJ, Witt CM, Linde K. Acupuncture for chronic pain: individual patient data meta-analysis. *Arch Intern Med*. 172:1444-1453, 2012
93. Vickers AJ, Cronin AM, Maschino AC, Lewith G, Macpherson H, Victor N, Sherman KJ, Witt C, Linde K. Individual patient data meta-analysis of acupuncture for chronic pain: protocol of the Acupuncture Trialists' Collaboration. *Trials* 11:90, 2010
94. Vickers AJ, Maschino AC, Lewith G, MacPherson H, Sherman KJ, Witt CM, Acupuncture Trialists C. Responses to the Acupuncture Trialists' Collaboration individual patient data meta-analysis. *Acupunct Med*. 31:98-100, 2013
95. Vickers AJ, Rees RW, Zollman CE, McCarney R, Smith CM, Ellis N, Fisher P, Van Haselen R. Acupuncture for chronic headache in primary care: large, pragmatic, randomised trial. *Bmj*. 328:744, 2004

96. Wallasch TM, Weinschuetz T, Mueller B, Kropp P. Cerebrovascular response in migraineurs during prophylactic treatment with acupuncture: a randomized controlled trial. *J Altern Complement Med.* 18:777-783, 2012
97. Wand BM, Abbaszadeh S, Smith AJ, Catley MJ, Moseley GL. Acupuncture applied as a sensory discrimination training tool decreases movement-related pain in patients with chronic low back pain more than acupuncture alone: a randomised cross-over experiment. *British journal of sports medicine.* 47:1085-1089, 2013
98. Wang LP, Zhang XZ, Guo J, Liu HL, Zhang Y, Liu CZ, Yi JH, Wang LP, Zhao JP, Li SS. Efficacy of acupuncture for migraine prophylaxis: a single-blinded, double-dummy, randomized controlled trial. *Pain.* 152:1864-1871, 2011
99. Wang LP, Zhang XZ, Guo J, Liu HL, Zhang Y, Liu CZ, Yi JH, Wang LP, Zhao JP, Li SS. Efficacy of acupuncture for acute migraine attack: a multicenter single blinded, randomized controlled trial. *Pain medicine (Malden, Mass.).* 13:623-630, 2012
100. Wang Y, Xue CC. Acupuncture for Frequent Migraine: A Randomized, Patient/Assessor Blinded, Controlled Trial with One-Year Follow-Up. 2015:920353, 2015
101. Weiner DK, Moore CG, Morone NE, Lee ES, Kent Kwoh C. Efficacy of periosteal stimulation for chronic pain associated with advanced knee osteoarthritis: a randomized, controlled clinical trial. *Clin Ther.* 35:1703-1720.e1705, 2013
102. Weiss J, Quante S, Xue F, Muche R, Reuss-Borst M. Effectiveness and acceptance of acupuncture in patients with chronic low back pain: results of a prospective, randomized, controlled trial. *J Altern Complement Med.* 19:935-941, 2013
103. White P, Bishop FL, Prescott P, Scott C, Little P, Lewith G. Practice, practitioner, or placebo? A multifactorial, mixed-methods randomized controlled trial of acupuncture. *Pain.* 153:455-462, 2012
104. White P, Lewith G, Prescott P, Conway J. Acupuncture versus placebo for the treatment of chronic mechanical neck pain: a randomized, controlled trial. *Ann Intern Med.* 141:911-919, 2004
105. Whitehurst DG, Bryan S, Hay EM, Thomas E, Young J, Foster NE. Cost-effectiveness of acupuncture care as an adjunct to exercise-based physical therapy for osteoarthritis of the knee. *Physical therapy.* 91:630-641, 2011
106. Wilke J, Vogt L, Niederer D, Hubscher M, Rothmayr J, Ivkovic D, Rickert M, Banzer W. Short-term effects of acupuncture and stretching on myofascial trigger point pain of the neck: a blinded, placebo-controlled RCT. *Complement Ther Med.* 22:835-841, 2014
107. Williamson L, Wyatt MR, Yein K, Melton JT. Severe knee osteoarthritis: a randomized controlled trial of acupuncture, physiotherapy (supervised exercise) and standard management for patients awaiting knee replacement. *Rheumatology (Oxford).* 46:1445-1449, 2007
108. Witt C, Brinkhaus B, Jena S, Linde K, Streng A, Wagenpfeil S, Hummelsberger J, Walther HU, Melchart D, Willich SN. Acupuncture in patients with osteoarthritis of the knee: a randomised trial. *Lancet.* 366:136-143, 2005
109. Witt CM, Jena S, Brinkhaus B, Liecker B, Wegscheider K, Willich SN. Acupuncture for patients with chronic neck pain. *Pain.* 2006
110. Witt CM, Jena S, Brinkhaus B, Liecker B, Wegscheider K, Willich SN. Acupuncture in patients with osteoarthritis of the knee or hip: a randomized, controlled trial with an additional nonrandomized arm. *Arthritis Rheum.* 54:3485-3493, 2006
111. Witt CM, Jena S, Selim D, Brinkhaus B, Reinhold T, Wruck K, Liecker B, Linde K, Wegscheider K, Willich SN. Pragmatic Randomized Trial Evaluating the Clinical and Economic Effectiveness of Acupuncture for Chronic Low Back Pain. *Am J Epidemiol.* 164:487-496, 2006
112. Witt CM, Schutzler L, Ludtke R, Wegscheider K, Willich SN. Patient characteristics and variation in treatment outcomes: which patients benefit most from acupuncture for chronic pain? *Clin J Pain.* 27:550-555, 2011
113. Yang CP, Chang MH, Li TC, Hsieh CL, Hwang KL, Chang HH. Predicting prognostic factors in a randomized controlled trial of acupuncture versus topiramate treatment in patients with chronic migraine. *Clin J Pain.* 29:982-987, 2013
114. Yang CP, Chang MH, Liu PE, Li TC, Hsieh CL, Hwang KL, Chang HH. Acupuncture versus topiramate in chronic migraine prophylaxis: a randomized clinical trial. *Cephalalgia.* 31:1510-1521, 2011
115. Yang J, Zeng F, Feng Y, Fang L, Qin W, Liu X, Song W, Xie H, Chen J, Liang F. A PET-CT study on the specificity of acupoints through acupuncture treatment in migraine patients. *BMC complementary and alternative medicine.* 12:123, 2012

116. Yang M, Yang J, Zeng F, Liu P, Lai Z, Deng S, Fang L, Song W, Xie H, Liang F. Electroacupuncture stimulation at sub-specific acupoint and non-acupoint induced distinct brain glucose metabolism change in migraineurs: a PET-CT study. *Journal of translational medicine*. 12:351, 2014
117. Yun M, Shao Y, Zhang Y, He S, Xiong N, Zhang J, Guo M, Liu D, Luo Y, Guo L, Yan J. Hegu acupuncture for chronic low-back pain: a randomized controlled trial. *J Altern Complement Med*. 18:130-136, 2012
118. Zhang SP, Chiu TT, Chiu SN. Long-term efficacy of electroacupuncture for chronic neck pain: a randomised controlled trial. *Hong Kong medical journal = Xianggang yi xue za zhi / Hong Kong Academy of Medicine*. 19 Suppl 9:36-39, 2013
119. Zhang WT, Jin Z, Cui GH, Zhang KL, Zhang L, Zeng YW, Luo F, Chen AC, Han JS. Relations between brain network activation and analgesic effect induced by low vs. high frequency electrical acupoint stimulation in different subjects: a functional magnetic resonance imaging study. *Brain Res*. 982:168-178, 2003
120. Zhao H, Nie W, Sun Y, Li S, Yang S, Meng F, Zhang L, Wang F, Huang S. Warm Needling Therapy and Acupuncture at Meridian-Sinew Sites Based on the Meridian-Sinew Theory: Hemiplegic Shoulder Pain. *Evid Based Complement Alternat Med*. 2015:694973, 2015
121. Zhao L, Liu J, Zhang F, Dong X, Peng Y, Qin W, Wu F, Li Y, Yuan K, von Deneen KM, Gong Q, Tang Z, Liang F. Effects of long-term acupuncture treatment on resting-state brain activity in migraine patients: a randomized controlled trial on active acupoints and inactive acupoints. *PloS one*. 9:e99538, 2014

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.

Figure 2. Forest plots for the comparison of acupuncture with no-acupuncture control. There were fewer than 3 trials for shoulder pain, so no meta-analyses were performed. Weights reported are fixed-effects weights calculated using inverse-variance weighting.

Figure 3. Forest plots for the comparison of true and sham acupuncture. Weights reported are fixed-effects weights calculated using inverse-variance weighting.

Figure 4. Forest plot showing the difference in pain change scores between acupuncture and no acupuncture control groups (a) and between acupuncture and sham acupuncture groups (b) over time. A coefficient of 0.01 means that the difference between acupuncture and control increases by 0.01 standard deviations for each 3 months following the end of treatment.

Supplementary Materials

Figure S1. Forest plots for the comparison of acupuncture with no-acupuncture control. There were fewer than 3 trials for shoulder pain, so no meta-analyses were performed. Weights reported are random-effects weights calculated using the DerSimonian and Laird method.

Figure S2. Forest plots for the comparison of true and sham acupuncture. Weights reported are random-effects weights calculated using the DerSimonian and Laird method.

Table 1. Characteristics of included studies

Indication (n=44)	Pain Type	Control Group	Primary Outcome Measure	Time Point
Chronic headache (n=9)	Migraine (n=3) ^{26, 59, 63} , tension-type headache (n=3) ^{23, 28, 71} , both ^{31, 43, 95} (n=3)	Sham control (n=5) ^{26, 28, 59, 63, 71} No acupuncture control (n=7); ancillary care (n=2) ^{23, 31} ; usual care (n=4) ^{43, 63, 71, 95} ; guideline care (n=1) ²⁶	Severity score (n=2) ^{23, 95} ; days with headache (n=3) ^{28, 43, 71} ; days with migraine (n=2) ^{26, 59} ; days with moderate-to-severe pain (n=1) ⁶³ ; Migraine Disability Assessment (MIDAS) (n=1) ³¹	1 mo (n=1) ²³ 2 mo (n=1) ³¹ 3 mo (n=3) ^{43, 63, 71} 4 mo (n=1) ⁵⁹ 6 mo (n=2) ^{26, 28} 12 mo (n=1) ⁹⁵
Nonspecific musculoskeletal pain (back and neck) (n=18)	Back (n=12) ^{11, 13, 18, 19, 36, 40, 48, 49, 74, 87, 102, 111} ; neck (n=6) ^{41, 67, 79, 91, 104, 109}	Sham control (n=10) ^{11, 13, 19, 36, 41, 48, 49, 74, 91, 104} No acupuncture control (n=12); Ancillary care (n=3) ^{40, 74, 102} ; usual care (n=7) ^{11, 19, 67, 79, 87, 109, 111} ; non-specific advice (n=1) ¹⁸ ; guideline care (n=1) ³⁶	VAS (n=7) ^{11, 13, 41, 49, 74, 91, 104} ; Roland Morris Disability Questionnaire (n=3) ^{18, 19, 48} ; Northwick Park Neck Pain Questionnaire (n=2) ^{67, 79} ; SF-36 Bodily pain (n=2) ^{87, 102} ; Hannover Functional Questionnaire (n=1) ¹¹¹ ; Von Korff pain score (n=1) ³⁶ ; Oswestry Disability Index (n=1) ⁴⁰	1 mo (n=4) ^{41, 49, 91, 104} 2 mo (n=3) ^{11, 18, 19} 3 mo (n=5) ^{48, 74, 79, 109, 111} 4 mo (n=1) ¹⁰² 6 mo (n=2) ^{36, 40} 8 mo (n=1) ¹³ 12 mo (n=1) ⁶⁷ 24 mo (n=1) ⁸⁷
Osteoarthritis (n=13)		Sham control (n=10) ^{8, 16, 33, 39, 70, 80, 85, 89, 103, 108} No acupuncture control (n=10); ancillary care (n=3) ^{33, 70, 80} ; usual care (n=5) ^{39, 56, 85, 108, 110} ; nonspecific advice (n=2) ^{8, 107}	WOMAC (n=5) ^{16, 56, 70, 108, 110} ; WOMAC Pain subscore (n=4) ^{8, 33, 80, 89} ; Oxford Knee score questionnaire (n=1) ¹⁰⁷ ; VAS ¹⁰³ (n=1); knee pain (0-10) (n=1) ³⁹ ; Joint-specific Multidimensional Assessment of Pain (n=1) ⁸⁵	1 mo (n=1) ¹⁰³ 2 mo (n=3) ^{70, 107, 108} 3 mo (n=6) ^{16, 39, 56, 85, 89, 110} 6 mo (n=3) ^{8, 33, 80}
Shoulder pain (n=4)		Sham control (n=4) ^{35, 52, 75, 90} No-acupuncture control (n=1); ancillary care (n=1) ⁷⁵	Constant-Murley score (n=2) ^{52, 90} ; VAS (n=2) ^{35, 75}	1 mo (n=2) ^{52, 90} 6 mo (n=2) ^{35, 75}

Table 2. Primary Analyses, N=44 trials. Acupuncture is superior to control at $p<0.001$ except where indicated

		Sham				No acupuncture control			
Analysis	Indication	No. of studies	FE (95% CI)	Heterogeneity p-value	RE (95% CI)	No. of studies	FE (95% CI)	Heterogeneity p-value	RE (95% CI)
Main Analysis	Non-specific musculoskeletal pain	10	0.30 (0.21, 0.38)	$p<0.001$	0.49 (0.16, 0.81)	12	0.54 (0.50, 0.57)	$p<0.001$	0.50 (0.38, 0.63)
	Osteoarthritis	9	0.24 (0.17, 0.31)	$p<0.001$	0.45 (0.15, 0.75)	10	0.63 (0.56, 0.69)	$p<0.001$	0.74 (0.46, 1.01)
	Chronic headache	5	0.16 (0.08, 0.25)	$p=0.4$	0.16 (0.08, 0.25)	7	0.44 (0.39, 0.48)	$p<0.001$	0.56 (0.35, 0.76)
	Shoulder	4	0.57 (0.44, 0.69)	$p=0.4$	0.57 (0.44, 0.69)	0	No trials		
Exclusion of Vas trials	Non-specific musculoskeletal pain	9	0.19 (0.11, 0.28)	$p=0.001$	0.31 (0.13, 0.48)				
	Osteoarthritis	8	0.18 (0.10, 0.25)	$p<0.001$	0.35 (0.07, 0.62)				
	Shoulder	3	0.58 (0.42, 0.74)	$p=0.2$	0.61 (0.40, 0.81)				
Separate pain types	Back pain	7	0.17 (0.07, 0.26)	$p<0.001$	0.30 (0.08, 0.52)	9	0.46 (0.41, 0.50)	$p<0.001$	0.52 (0.37, 0.67)
	Neck pain	3	0.83 (0.64, 1.01)	$p<0.001$	0.82 (-0.11, 1.75)				
Excluding trials with summary data only	Non-specific musculoskeletal pain	9	0.27 (0.19, 0.35)	$p<0.001$	0.44 (0.11, 0.78)	11	0.53 (0.50, 0.56)	$p<0.001$	0.45 (0.33, 0.57)
	Osteoarthritis	8	0.19 (0.12, 0.26)	$p<0.001$	0.26 (0.04, 0.48)	9	0.59 (0.52, 0.65)	$p<0.001$	0.59 (0.37, 0.82)
	Chronic headache					5	0.43 (0.38, 0.47)	$p<0.001$	0.44 (0.24, 0.64)
	Shoulder	3	0.62 (0.46, 0.77)	$p=0.4$	0.62 (0.46, 0.77)				
Excluding trials with possible bias due to blinding	Non-specific musculoskeletal pain	7	0.28 (0.19, 0.37)	$p<0.001$	0.51 (0.09, 0.93)				
	Osteoarthritis	9	0.23 (0.16, 0.31)	$p<0.001$	0.44 (0.13, 0.75)				
	Chronic headache*	3	0.15 (0.03, 0.26)	$p=0.15$	0.12 (-0.05, 0.29)				
Including trials with high likelihood of bias due to blinding	Osteoarthritis	10	0.23 (0.17, 0.30)	$p<0.001$	0.42 (0.14, 0.70)				
Multiple imputation	Non-specific musculoskeletal pain	10	0.29 (0.21, 0.37)	$p<0.001$	0.49 (0.16, 0.81)	12	0.54 (0.50, 0.57)	$p<0.001$	0.51 (0.38, 0.63)
	Osteoarthritis	9	0.24 (0.17, 0.31)	$p<0.001$	0.45 (0.15, 0.75)	10	0.63 (0.57, 0.70)	$p<0.001$	0.74 (0.46, 1.01)
	Chronic headache	5	0.17 (0.08, 0.25)	$p=0.4$	0.17 (0.08, 0.25)	7	0.44 (0.40, 0.49)	$p<0.001$	0.55 (0.35, 0.75)
	Shoulder	4	0.56 (0.44, 0.69)	$p=0.4$	0.56 (0.44, 0.69)				
Excluding trials where both acupuncture and control groups received additional treatments	Non-specific musculoskeletal pain					10	0.54 (0.51, 0.57)	$p<0.001$	0.54 (0.40, 0.67)
	Osteoarthritis	4	0.21 (0.11, 0.31)	$p=0.081$	0.22 (0.07, 0.38)	7	0.70 (0.62, 0.78)	$p<0.001$	0.70 (0.47, 0.93)
	Chronic headache					5	0.43 (0.38, 0.47)	$p<0.001$	0.44 (0.24, 0.64)
	Shoulder	3	0.58 (0.42, 0.74)	$p=0.2$	0.61 (0.40, 0.81)				

* $p=0.015$

Table 3. Sensitivity analyses including only pain endpoints measured between 2 and 3 months after randomization.

Analysis	Indication	Sham				No acupuncture control			
		No. of studies	FE (95% CI)	Heterogeneity p-value	RE (95% CI)	No. of studies	FE (95% CI)	Heterogeneity p-value	RE (95% CI)
Main Analysis	Non-specific musculoskeletal pain	5	0.13 (0.01, 0.25)	p=0.005	0.23 (-0.03, 0.49)	9	0.60 (0.56, 0.64)	p<0.0001	0.47 (0.34, 0.61)
	Osteoarthritis	7	0.31 (0.23, 0.39)	p<0.0001	0.69 (0.24, 1.14)	9	0.73 (0.66, 0.80)	p<0.0001	0.88 (0.61, 1.15)
	Chronic headache	5	0.14 (0.06, 0.22)	p=0.4	0.14 (0.06, 0.22)	7	0.43 (0.38, 0.47)	p<0.0001	0.45 (0.27, 0.63)
	Shoulder	2	No meta-analysis						

Table 4. Trial-level acupuncture characteristics, N=39. Counts for point prescription sum to 40 because one trial had two acupuncture groups, with each group receiving acupuncture based on a different point prescription.

Style of Acupuncture	
Combination of traditional Chinese and Western	9 (23%)
Traditional Chinese techniques	23 (59%)
Western	7 (18%)
Point Prescription	
Fixed needle formula	9 (23%)
Flexible formula	18 (45%)
Individualized	13 (33%)
Location of needles	
Both Local and Distal Points	37 (95%)
Distal Points Only	2 (5.1%)
Electrical stimulation allowed	11 (28%)
Manual stimulation allowed	36 (92%)
Moxibustion allowed	6 (15%)
Other Adjunctive Therapies Allowed	8 (21%)
De Qi attempted (N=35)	33 (94%)
Acupuncture-specific patient practitioner interactions	16 (40%)
Minimum years of experience required	
No requirement specified (0 years)	14 (36%)
6 months to 2 years	7 (18%)
3-4 years	13 (33%)
5-9 years	3 (7.7%)
10 years	2 (5.1%)
Maximum number of sessions	
1-5	3 (7.7%)
6-10	19 (49%)
11-15	12 (31%)
16-20	1 (2.6%)
21-25	2 (5.1%)
26-30	2 (5.1%)
Frequency of sessions (mean number of sessions per week)	
0.88	1 (2.6%)
1	19 (49%)
1.43	1 (2.6%)
1.5	7 (18%)
1.67	1 (2.6%)
2	9 (23%)
5	1 (2.6%)
Mean duration of sessions, rounded to whole numbers (N=34)	
15-19 minutes	1 (2.9%)
20-24 minutes	11 (32%)
25-29 minutes	6 (18%)
30+ minutes	16 (47%)
Mean number of needles used (N=33)	
1-4	3 (9.1%)
5-9	11 (33%)

10-14	12 (36%)
15-20	7 (21%)

Accepted Manuscript

Table 5. Patient-level acupuncture characteristics, n=20,827.

Number of Sessions	
0	441 (2.1%)
1-5	515 (2.5%)
6-10	8003 (38%)
11-15	2065 (10%)
16-20	40 (0.2%)
21-30	15 (<0.1%)
Missing	1989 (10%)
Not reported	7759 (37%)
Average Session Duration	
2-15	163 (0.8%)
15-30	2668 (13%)
31-45	377 (1.8%)
46-60	25 (0.1%)
60+	1 (<0.1%)
Missing	896 (4.3%)
Not reported	16697 (80%)
Average Number of Needles	
2-5	22 (0.1%)
6-10	910 (4.4%)
11-15	762 (3.7%)
16-20	825 (4.0%)
21-25	199 (1.0%)
26+	30 (0.1%)
Missing	1621 (7.8%)
Not reported	16458 (79%)
Age of Physician/Acupuncturist	
30-35	298 (1.4%)
36-40	2119 (10%)
41-45	2630 (13%)
46-50	2407 (12%)
51-55	1701 (8.2%)
56-60	872 (4.2%)
60+	303 (1.5%)
Missing	368 (1.8%)
Not reported	10129 (49%)
Physician/Acupuncturist Sex	
Female	3626 (17%)
Male	7002 (34%)
Missing	70 (0.3%)
Not reported	10129 (49%)

Table 6. Results of univariate meta-regression analyses for the effect of acupuncture characteristics on acupuncture effect. β is an estimate of the change in the effect of acupuncture in terms of standardized difference compared to controls for each characteristic; a positive β indicates a larger effect of acupuncture compared to controls for trials. N is number of trials. The number of patients in the analysis and number of patients in included trials are given in parentheses where applicable.

	Sham acupuncture				No acupuncture control			
	N	β	95% CI	p value	N	β	95% CI	p value
Style of acupuncture	25				25			
Some TCM vs. Western only		-0.00	-0.49, 0.48	>0.9		0.10	-0.55, 0.74	0.8
TCM only vs. some Western		0.02	-0.38, 0.42	0.9		-0.07	-0.42, 0.28	0.7
Point prescription	25				25			
Fixed needle formula		Ref.		0.6		Ref.		0.075
Flexible formula		0.20	-0.21, 0.60			0.01	-0.45, 0.46	
Fully individualized		-0.01	-0.75, 0.73			-0.34	-0.79, 0.10	
Electrical stimulation allowed	25	0.32	-0.11, 0.75	0.14	25	-0.12	-0.50, 0.26	0.5
Manual stimulation allowed	25	0.26	-0.42, 0.95	0.5	25	-0.38	-0.99, 0.23	0.2
Moxibustion allowed		No trials allowed			25	-0.32	-0.71, 0.06	0.10
Other adjunctive treatment allowed	25	-0.04	-1.00, 0.92	0.9	25	-0.22	-0.59, 0.16	0.3
De qi attempted	25	0.29	-0.67, 1.24	0.6	21	0.74	-0.04, 1.52	0.063
Acupuncture-specific patient practitioner interactions allowed	25	-0.03	-0.50, 0.44	0.9	25	-0.05	-0.38, 0.28	0.8
Minimum experience required (years)	25	0.04	-0.05, 0.13	0.4	25	0.05	-0.03, 0.12	0.2
Maximum number of sessions (per 5 sessions)	25	-0.01	-0.23, 0.22	0.9	25	0.01	-0.12, 0.14	0.9
Patient-level analysis	5 (1317/ 1377)	0.09	-0.31, 0.48	0.7	5 (8036/ 10157)	0.10	-0.01, 0.21	0.001
Patient-level analysis, including Hinman trial	6 (1421/1517)	-0.03	-0.36, -0.30	0.9				
Frequency of sessions (per week)	25	-0.06	-0.29, 0.18	0.6	25	0.21	-0.22, 0.64	0.3
Duration of sessions (per 5 minutes)	25	0.06	-0.13, 0.25	0.5	20	-0.06	-0.25, 0.13	0.5
Patient-level analysis	6 (2863/2969)	0.01	-0.08, 0.09	0.9				
Number of needles used (per 5 needles)	25	0.05	-0.17, 0.27	0.6	19	0.16	-0.05, 0.38	0.13
Patient-level analysis	5 (2232/2317)	0.04	-0.08, 0.16	0.5				
Age of practitioner (per 5 years)								
Patient-level analysis					6 (9127/10550)	-0.01	-0.04, 0.02	0.5
Male practitioner								
Patient-level analysis					6 (9384/10550)	-0.07	-0.16, 0.02	0.084

Table 7. Differences in effect size (in SD) between acupuncture and sham acupuncture groups (N=25) and between acupuncture and no acupuncture control groups (N=24). Total number of sham acupuncture-controlled trials sums to 26 because one trial had two different types of sham acupuncture control.

Sham Acupuncture			
Type of Control Group	N	Effect Size (95% CI)	p value
Penetrating needle sham	11	0.17 (0.11, 0.22)	<0.0001
Excluding B blinding grades	9	0.16 (0.09, 0.24)	<0.0001
Non-penetrating needle and non-needle sham	15	0.48 (0.22, 0.74)	0.0003
Excluding B blinding grades	11	0.51 (0.16, 0.86)	0.004
Including Hinman trial	16	0.46 (0.21, 0.70)	0.0003
Excluding Vas trials	12	0.27 (0.10, 0.44)	0.002
Non-penetrating needle sham	10	0.52 (0.14, 0.91)	0.007
Excluding Vas trials	7	0.22 (-0.05, 0.49)	0.11
Non-needle sham	5	0.37 (0.21, 0.52)	<0.0001
Including Hinman trial	6	0.32 (0.18, 0.46)	<0.0001
True acupuncture points (no penetrating needle sham)	12	0.48 (0.15, 0.80)	0.004
Excluding B blinding grades	10	0.51 (0.12, 0.89)	0.010
Including Hinman trial	13	0.45 (0.15, 0.75)	0.003
Excluding Vas trials	10	0.25 (0.06, 0.44)	0.011
Non-acupuncture points (no penetrating needle sham)	3	0.52 (0.35, 0.69)	<0.0001
Excluding Vas trials	2	0.47 (0.13, 0.81)	0.007
No Acupuncture Control			
Type of Control Group	N	Effect Size (95% CI)	p value
High intensity	5	0.34 (0.11, 0.57)	0.003
Usual care and low intensity	19	0.56 (0.43, 0.69)	<0.0001
Usual care	17	0.50 (0.40, 0.60)	<0.0001
Low intensity	2	1.14 (0.71, 1.58)	<0.0001

Table 8. Differences in effect size between different types of control group. A negative effect size indicates that there is a smaller difference in effect between acupuncture and control for group 1 than for group 2, that is, the effect of control group 1 is more similar to verum acupuncture than the effect of control group 2.

Sham Acupuncture			
Group 1	Group 2	Effect Size (95% CI)	p value
Penetrating needle sham	Non-penetrating and non-needle sham	-0.30 (-0.60, -0.00)	0.047
Excluding B blinding grades		-0.33 (-0.72, 0.05)	0.088
Including Hinman trial		-0.28 (-0.57, 0.01)	0.061
Excluding Vas trials		-0.07 (-0.24, 0.10)	0.4
Non-penetrating needle sham	Non-needle sham	0.13 (-0.44, 0.70)	0.6
Including Hinman trial		0.18 (-0.34, 0.70)	0.5
Excluding Vas trials		-0.18 (-0.52, 0.17)	0.3
True acupuncture points, excluding penetrating needle sham	Non-acupuncture points, excluding penetrating needle sham	-0.02 (-0.70, 0.66)	0.9
Including Hinman trial		-0.05 (-0.71, 0.61)	0.9
Excluding Vas trials		-0.22 (-0.75, 0.30)	0.4
No Acupuncture Controls			
Group 1	Group 2	Effect Size (95% CI)	p value
High intensity	Usual care and low intensity	-0.23 (-0.50, 0.05)	0.11
High intensity	Low intensity	-0.81 (-1.26, -0.36)	0.0004
Usual care	Low intensity	-0.65 (-0.98, -0.31)	0.0002

Table 9. Trials with sham and no acupuncture control and time points assessed after the end of treatment

Trial Name	Pain Condition	Average Length of Treatment	Sham Acupuncture		No acupuncture control		
			Time Points after End of Treatment	Included in meta-analysis	Control patients offered acupuncture treatment (Crossover)	Time Points after End of Treatment	Included in meta-analysis
Carlsson 2001 ¹³	Low Back Pain	8 weeks	Weeks 5 and 18	Yes			
Chen 2013 ¹⁶	Osteoarthritis	12 weeks	End of treatment and week 14	Yes			
Endres 2007 ²⁸	Headache	6 weeks	End of treatment and weeks 7 and 20	Yes			
Guerra de Hoyos 2004 ³⁵	Shoulder	8 weeks	Weeks 5 and 18	Yes			
Irnich 2001 ⁴¹	Neck	3 weeks	Weeks 1 and 10	Yes			
Kennedy 2008 ⁴⁸	Low Back Pain	5 weeks	End of treatment and week 7	Yes			
Kerr 2003 ⁴⁹	Low Back Pain	6 weeks	None	No			
Kleinhenz 1999 ⁵²	Shoulder	4 weeks	End of treatment	No			
Li 2012 ⁵⁹	Migraine	4 weeks	End of treatment and week 4	Yes			
Vas 2004 ⁸⁹	Osteoarthritis	12 weeks	Week 1	No			
Vas 2006 ⁹¹	Neck	3 weeks	Weeks 1 and 25	Yes			
Vas 2008 ⁹⁰	Shoulder	3 weeks	Weeks 1 and 10	Yes			
White 2004 ¹⁰⁴	Neck	4 weeks	End of treatment and weeks 1 through 8	Yes			
White 2012 ¹⁰³	Osteoarthritis	4 weeks	End of treatment and week 1	Yes			
Berman 2004 ⁸	Osteoarthritis	26 weeks	End of treatment	No	No	End of treatment	No
Brinkhaus 2006 ¹¹	Low Back Pain	8 weeks	End of treatment and weeks 18 and 44	Yes	At 8 weeks	End of treatment	No
Cherkin 2009 ¹⁹	Low Back Pain	7 weeks	Weeks 1, 19 and 45	Yes	No	Weeks 1, 19 and 45	Yes
Diener 2006 ²⁶	Migraine	6 weeks	End of treatment and weeks 7 and 20	Yes	No	End of treatment and weeks 7 and 20	Yes
Foster 2007 ³³	Osteoarthritis	3 weeks	Weeks 3, 23 and 49	Yes	No	Weeks 3, 23 and 49	Yes
Haake 2007 ³⁶	Low Back Pain	6 weeks	End of treatment and weeks 7 and 20	Yes	No	End of treatment and weeks 7 and 20	Yes
Linde 2005 ⁶³	Migraine	8 weeks	End of treatment and weeks 4 and 16	Yes	At 12 weeks	Week 4	No
Melchart 2005 ⁷¹	Headache	8 weeks	End of treatment and weeks 4 and 16	Yes	At 12 weeks	Week 4	No
Scharf 2006 ⁸⁰	Osteoarthritis	6 weeks	Weeks 7 and 20	Yes	No	Weeks 7 and 20	Yes
Suarez-Almazor 2010 ⁸⁵	Osteoarthritis	6 weeks	End of treatment and week 7	Yes	No	Week 7	No
Witt 2005 ¹⁰⁸	Osteoarthritis	8 weeks	End of treatment and weeks 18 and 44	Yes	At 8 weeks	End of treatment	No

Cherkin 2001 ¹⁸	Low Back Pain	10 weeks			No	End of treatment and week 42	Yes
Hinman 2014 ³⁹	Osteoarthritis	12 weeks			No	End of treatment and week 40	Yes
Hunter 2012 ⁴⁰	Low Back Pain	6 weeks			No	Weeks 2, 7 and 20	Yes
Jena 2008 ⁴³	Headache	12 weeks			At 12 weeks	All measurements after crossover	No
Lansdown 2009 ⁵⁶	Osteoarthritis	10 weeks			No	Weeks 3 and 42	Yes
MacPherson 2015 ⁶⁷	Neck	16 weeks			No	Weeks 10 and 36	Yes
Thomas 2006 ⁸⁷	Low Back Pain	12 weeks			No	Weeks 1, 40 and 92	Yes
Salter 2006 ⁷⁹	Neck	12 weeks			No	Week 1	No
Vickers 2004 ⁹⁵	Headache	6 weeks			No	Weeks 1 and 40	Yes
Weiss 2013 ¹⁰²	Low Back Pain	4 weeks			No	End of treatment and week 13	Yes
Williamson 2007 ¹⁰⁷	Osteoarthritis	6 weeks			No	Weeks 1 and 6	Yes
Witt 2006 ¹⁰⁹	Neck	12 weeks			At 12 weeks	All measurements after crossover	No
Witt 2006 ¹¹⁰	Osteoarthritis	12 weeks			At 12 weeks	All measurements after crossover	No
Witt ARC 2006 ¹¹¹	Low Back Pain	12 weeks			At 12 weeks	All measurements after crossover	No






Figure 1 2017-10-24_bestsetConverted.png

Accepted Manuscript

Osteoarthritis



Headache



Musculoskeletal Pain






Figure 2 No Acupuncture Control.eps

Accepted Manuscript


Osteoarthritis



Headache



Musculoskeletal Pain



Specific Shoulder Pain




Figure 3 Sham Control.eps

Accepted Manuscript




Figure 4a Time Course No Acupuncture.eps

Accepted Manuscript






Figure 4b Time Course Sham.eps

Accepted Manuscript

Osteoarthritis



Headache



Musculoskeletal Pain






Figure S1 No Acupuncture Control RE.eps

Accepted Manuscript


Osteoarthritis



Headache



Musculoskeletal Pain



Specific Shoulder Pain




Figure S2 Sham Control RE.eps

Accepted Manuscript